

# Understanding how effective 'induced after-death communication' therapy is in helping people deal with grief and what factors influence the results

<b>Submission date</b> 23/03/2025	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol <input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results <input type="checkbox"/> Individual participant data
<b>Registration date</b> 24/03/2025	<b>Overall study status</b> Completed	
<b>Last Edited</b> 04/02/2026	<b>Condition category</b> Other	

## Plain English summary of protocol

### Background and study aims

Grief is a normal human experience, but some people experience intense sadness that lasts for months or years, making everyday life difficult. Induced After-Death Communication (IADC) therapy is a brief, two-session treatment that may help people process their grief and feel reconnected with loved ones who have passed away. This study aimed to test whether IADC therapy helps reduce grief symptoms, and to explore which factors might predict better outcomes.

### Who can participate?

Adults aged 18 and older who were grieving the loss of a loved one and had experienced that loss at least six months prior were eligible. All participants needed to speak English and be able to tolerate strong emotions during therapy.

### What does the study involve?

Participants first completed a screening interview to ensure suitability. They then completed online surveys at four points: one month before therapy, just before therapy, just after therapy, and one month after therapy. Therapy involved two 90-minute sessions focused on reducing the sadness of grief using a specialised technique involving eye movements or tapping (bilateral stimulation). Some participants also reported sensing the presence of their deceased loved one during therapy.

### What are the possible benefits and risks of participating?

Many participants reported feeling "lighter" and better able to move forward after the therapy. Benefits may include reduced sadness and stronger feelings of connection with the deceased. Risks include temporary emotional discomfort when recalling sad memories, and feeling tired after therapy sessions.

**Where is the study run from?**

The study was conducted by two psychologists in private practice—one in Adelaide, South Australia, and one in Queensland. Some sessions were delivered in person, others via Zoom.

**When is the study starting and how long is it expected to run for?**

The study began in July 2023 and data collection finished in early 2024. The write-up is currently underway.

**Who is funding the study?**

This study received no external funding and was conducted voluntarily by the lead researcher.

**Who is the main contact?**

Dr Tom Nehmy

School of Psychology

University of Adelaide

[tom.nehmy@adelaide.edu.au](mailto:tom.nehmy@adelaide.edu.au)

## Contact information

**Type(s)**

Public, Scientific, Principal investigator

**Contact name**

Dr Thomas Nehmy

**ORCID ID**

<https://orcid.org/0009-0001-4606-948X>

**Contact details**

School of Psychology

The University of Adelaide

North Terrace Campus

Adelaide

Australia

5005

+61 8 8313 4763

[tom.nehmy@adelaide.edu.au](mailto:tom.nehmy@adelaide.edu.au)

## Additional identifiers

**Clinical Trials Information System (CTIS)**

Nil known

**ClinicalTrials.gov (NCT)**

Nil known

**Protocol serial number**

H-2023-207

## Study information

**Scientific Title**

Efficacy and predictors of outcome of Induced After-Death Communication (IADC) therapy in treating grief

**Acronym**

IADC Study

**Study objectives**

The primary hypothesis is that IADC Therapy will have therapeutic effects on grief symptoms and sadness.

**Ethics approval required**

Ethics approval required

**Ethics approval(s)**

approved 29/08/2023, Human Research Ethics Committee, University of Adelaide (LEVEL 3, RUNDLE MALL PLAZA 50 RUNDLE MALL, Adelaide, 5000, Australia; +61883135137; hrec@adelaide.edu.au), ref: H-2023-207

**Study design**

Interventional wait-list controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Treatment of grief in bereaved adults

**Interventions**

Bereaved English-speaking adults in Australia were provided with Induced After-Death Communication (IADC) Therapy following a one-month waitlist (control) period.

**Intervention Type**

Behavioural

**Primary outcome(s)**

1. Grief Symptoms as measured using the International Prolonged Grief Disorder Scale at Time 1 (start of waitlist period), Time 2 (end of one-month baseline waitlist period), and Time 4 (one-month follow-up post-treatment).
2. Average Daily Sadness - subjective self-report on a 0-10 scale at Time 1 (start of waitlist period), Time 2 (end of one-month baseline waitlist period), and Time 4 (one-month follow-up post-treatment).

**Key secondary outcome(s)**

1. Depression as measured by the Depression, Anxiety & Stress Scale (DASS-21)
2. Occurrence of after-death communication (ADC) as measured by self-report at Time 3 (immediately posttreatment) and Time 4 (one month follow-up)

3. Ongoing bonds as measured by the Continuing Presence subscale of the Ongoing Attachment Inventory at Time 1 (start of waitlist period), Time 2 (end of one-month baseline waitlist period), and Time 4 (one-month follow-up post-treatment).
4. Grief as measured by the Hogan Grief Reaction Checklist at Time 1 (start of waitlist period), Time 2 (end of one-month baseline waitlist period), and Time 4 (one-month follow-up post-treatment).

**Completion date**

16/07/2024

## Eligibility

**Key inclusion criteria**

1. English-speaking
2. Adults (>18 years)
3. Within Australia

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Mixed

**Lower age limit**

18 years

**Upper age limit**

110 years

**Sex**

All

**Total final enrolment**

45

**Key exclusion criteria**

1. Suicidality
2. If less than six months had passed since the loss
3. If clients were unwilling or unable to tolerate high levels of emotion or distress in confronting their grief and sadness.

**Date of first enrolment**

10/09/2023

**Date of final enrolment**

15/05/2024

# Locations

## Countries of recruitment

Australia

## Study participating centre

**Moveon Counselling**

168 Main Rd

Blackwood

Australia

5051

## Study participating centre

**John Daniels Psychology**

Suite 55, Level 2, Niecon Plaza, 19 Victoria Ave

Broadbeach

Australia

4218

## Study participating centre

**Healing Grief International Pty Ltd**

10 Scroop Rd

Hawthorndene

Australia

5051

# Sponsor information

## Organisation

The University of Adelaide

## ROR

<https://ror.org/00892tw58>

# Funder(s)

## Funder type

Other

**Funder Name**

Investigator initiated and funded

## Results and Publications

### Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not expected to be made available

#### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#"><u>Results article</u></a>		23/01/2026	04/02/2026	Yes	No
<a href="#"><u>Participant information sheet</u></a>	version 1.1	20/08/2023	24/03/2025	No	Yes