Assessing and training fear inhibition in spider phobia via eye-movements

Recruitment status No longer recruiting Overall study status Completed Condition category Mental and Behavioural Disorders	[X] Prospectively registered		
	[X] Protocol		
	[X] Statistical analysis plan [] Results		
	 Individual participant data 		
	[X] Record updated in last year		
	Io longer recruiting Overall study status completed Condition category		

Plain English summary of protocol

Background and study aim

Anxiety and fear can be characterized by a shift of attention in favor of threatening stimuli. According to the Attentional-Control-Theory, this bias is associated with a decreased inhibition of bottom-up perceptual processes. Inhibitory control can be investigated by means of the antisaccade task, in which participants are required to look into the opposite direction of a stimulus in the peripheral visual field. Higher antisaccade latencies and error rates thereby indicate inhibitory control deficits, which are considered a pathogenic factor in anxiety disorders. In line with this, previous studies have revealed an impaired antisaccade performance in participants with subclinical anxiety. However, studies investigating clinical populations are sparse.

This study has three main goals: First, we aim to compare the antisaccade performance of spider phobic patients with healthy control participants. Second, we aim to investigate associations between antisacccade performance and psychophysiological (heart rate, skin conductance level, startle response) as well as behavioral measures of fear (ratings, behavioral avoidance test) towards threatening stimuli. Third, this study aims to explore effects of an antisaccade training on inhibitory control (indexed by antisaccade performance), as well as psychophysiological and behavioral measures of fear.

Who can participate?

Patients with a specific phobia of the animal subtype (spiders) who fulfill the criteria for a specific phobia.

Healthy participants who do not fulfill the criteria for a specific phobia and do not exceed a value of 19 points in the spider phobia questionnaire (SPQ).

What does the study involve?

Participants take part in a brief screening and a diagnostic interview via telephone. If they meet the inclusion criteria, they will provide informed consent, fill in psychometric questionnaires and participate in laboratory assessments (2-3 hours for healthy controls and 3-4 hours for patients with spider phobia): First, a baseline assessment including a Behavioral Avoidance Test (BAT), a free viewing paradigm with physiological assessments (heart rate, skin conductance level, startle response), and the anti-saccade task with an eye-tracking assessment will be obtained. Second, participants are randomized into two intervention groups receiving an anti-saccade (experimental group) or a pro-saccade training (control group). Afterwards, the post-1assessment of the antisaccade task and the BAT are conducted.

From this point on the assessment continues only for participants with specific phobia. These switch training conditions and then participate in the post-2-assessment composed of the antisaccade task, the free viewing physiological assessment, and the BAT.

What are the possible benefits and risks of participating?

Results of this study shed light to pathogenic mechanisms involved in pathological anxiety and their modulation. As such, it may inform future research and clinical approaches of anxiety treatment. The risks of participating in the study are a short increase of anxiety, due to the anxiety-inducing stimuli, loud startle noises, and fatigue of the eyes.

Where is the study run from? University of Siegen (Germany) and University of Münster (Germany)

When is the study starting and how long is it expected to run for? June 2021 to December 2023

Who is funding the study? DGPs (German Society for Psychology: Biological Psychology and Neuropsychology) Movisens (Germany) Innovative Medizinische Forschung (IMF) of the medical faculty of Münster (Germany)

Who is the main contact? Dr. Kati Roesmann (Kati.Roesmann@uni-siegen.de) Fabian Breuer M.Sc. (breuerfa@uni-muenster.de)

Contact information

Type(s) Principal Investigator

Contact name Dr Kati Roesmann

ORCID ID http://orcid.org/0000-0001-6940-3551

Contact details Obergraben 23 Siegen Germany 57072 +49 271 740-5123 Kati.Roesmann@uni-siegen.de

Additional identifiers

EudraCT/CTIS number Nil known

IRAS number

ClinicalTrials.gov number Nil known

Secondary identifying numbers Nil known

Study information

Scientific Title

Inhibitory control and its modification in specific phobia - An antisaccade study

Acronym

SPIN

Study objectives

H1: Phobics will show higher latencies and error rates, i.e. a poorer performance, than healthy controls in the antisaccade paradigm.

H1.1: Differences in antisaccade performance between phobics and healthy controls are higher when using phobia-related vs. neutral stimuli.

H2: Latencies and error-rates in the anti-saccade task are correlated with physiological (skin conductance response, heart rate, startle-response) and behavioral measures of fear. H3: The anti-saccade training improves antisaccade performance (i.e., lower error rates, and shorter latencies)

H4: Changes in antisaccade performance are associated with changes in behavioral and physiological measures of fear.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 08/09/2021, Ethics Committee of the University of Siegen (Adolf-Reichwein-Str. 2a, NA, 57076, Germany; +49 0271 740-4819; ethikrat@uni-siegen.de), ref: ER_39/2021

Study design

Interventional randomized controlled trial

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) Other

Study type(s) Other

Participant information sheet

Not available in web format, please use contact information to request a participant information sheet

Health condition(s) or problem(s) studied

Inhibitory control in specific phobia of the animal typus (spider).

Interventions

Intervention group: Antisaccade training Control group: Prosaccade training Participants will be randomised into the two groups by a randomisation sheet.

Antisaccade Training: Visual stimuli (pictures of spiders) will be presented on a screen in the left or right peripheral visual field. Participants are instructed to look at the mirrored position on the screen. The duration of this training is 15 minutes including short breaks.

Prosaccade Training: Visual stimuli (pictures of neutral objects) will be presented on a screen in the left or right peripheral visual field. Participants are instructed to look at the presented stimulus. The duration of this training is 15 minutes including short breaks.

Intervention Type

Behavioural

Primary outcome measure

Antisaccade latencies will be measured in the antisaccade task at baseline, post-1-, and - for spider phobics only - post-2-assessment. All assessments take place on one day.

Secondary outcome measures

Antisaccade error rates will be measured in the antisaccade task at baseline, post-1-, and - for spider phobics only - post-2-assessment. All assessments take place on one day

Overall study start date

01/06/2021

Completion date

31/12/2023

Eligibility

Key inclusion criteria

- 1. 18-65 years
- 2. Specific phobia (animal-subtype: spider) and healthy controls
- 3. Normal or corrected-to normal vision
- 4. Normal hearing

Participant type(s) Mixed

Age group Adult

Lower age limit

18 Years

Upper age limit

65 Years

Sex

Both

Target number of participants

60 (30 patients with spider phobia, 30 healthy control participants)

Key exclusion criteria

1. Lifetime diagnosis of substance-related bipolar or psychotic disorder

2. Current psychiatric disorder or a psychiatric disorder in the past (exception for spider phobics: past mild to moderate depressive episode and/or current or past specific phobia of the animal typus)

- 3. Medication (Benzodiazepine, Barbiturate)
- 4. Neurological disorder (especially epilepsy)

5. Dementia

6. Injury of the central nervous system

7. Hearing disorder (also tinnitus anamnestic), subjective auditory hypersensitivity (like hyperacusis)

8. Regular nicotine consumption (>5 cigarettes/day)

9. Known allergy to bites of insects or arachnids

Date of first enrolment

01/03/2022

Date of final enrolment

31/12/2023

Locations

Countries of recruitment Germany

Study participating centre University of Siegen Obergraben 23 Siegen Germany 57072

Sponsor information

Organisation University of Siegen

Sponsor details Adolf-Reichwein-Straße 2 Siegen Germany 57076 +49 271 740-4106 tim.klucken@psychologie.uni-siegen.de

Sponsor type University/education

Website https://www.uni-siegen.de/start/

ROR https://ror.org/02azyry73

Funder(s)

Funder type University/education

Funder Name Medizinische Fakultät, Westfälische Wilhelms-Universität Münster

Alternative Name(s) Medical Faculty Münster, Medical Faculty, WWU Münster, Medizinische Fakultät Münster

Funding Body Type Government organisation

Funding Body Subtype Local government

Location Germany

Funder Name German Society for Psychology: Biological Psychology and Neuropsychology

Funder Name Movisens

Results and Publications

Publication and dissemination plan

Planned publication in a high-impact peer-reviewed journal

Intention to publish date

31/12/2024

Individual participant data (IPD) sharing plan

Data are available upon reasonable request. Kati.Roesmann@uni-siegen.de, breuerfa@uni-muenster.de

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient- facing?
<u>Statistical</u> Analysis Plan	A SAP on planned analyses of data obtained during the baseline assessment version 1.0	30/11 /2022	09/12 /2022	No	No
<u>Statistical</u> <u>Analysis Plan</u>	A SAP on planned analyses of data obtained during the intervention assessment version 1.0		12/04 /2023	No	No
Protocol article		19/12 /2023	08/01 /2024	Yes	No
<u>Statistical</u> Analysis Plan			02/09 /2024	No	No