

# A study in healthy smokers to determine the delivery of nicotine and satisfaction/craving relief gained from an e-cigarette containing different solutions (e-liquids)

<b>Submission date</b> 21/10/2014	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 04/11/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 07/06/2018	<b>Condition category</b> Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data

## Plain English Summary

### Background and study aims

Electronic cigarettes (e-cigarettes) are battery-powered vaporizers that have been developed to mimic the feel and experience of tobacco smoking. They work by vaporizing a liquid solution known as an e-liquid, which typically contains a mixture of propylene glycol, glycerin, nicotine and flavourings. There is some evidence to suggest that e-cigarettes may be safer than conventional tobacco smoking. Here, we want to compare the satisfaction and relief from craving nicotine people get from smoking one of three different e-cigarettes containing different solutions and a regular cigarette.

### Who can participate?

Adults aged 21-55, in good health and smoking at least 10 cigarettes a day for at least a year. Women are not allowed to take part if pregnant or breastfeeding.

### What does the study involve?

Once deemed eligible, participants attend the testing clinic four times during the study. During their first visit, they smoke a regular cigarette. Blood samples are then taken for nicotine analysis and the participants are asked to complete a questionnaire examining the effects of cigarette use on their cravings and urges to smoke. Their heart rate is monitored as they smoke and cigarette filters are collected for nicotine exposure analysis. After this visit, participants are given an e-cigarette to use for the next two days. They are randomly allocated one of three test e-cigarette solutions. After using the e-cigarette for two days they return to the clinic for the same nicotine analysis and cravings/urges analysis as before. Cartridges are weighed to estimate nicotine delivery from the e-cigarette. This process is repeated two more times so that each participant is given all three e-cigarette solutions to try. When at home, participants are asked to complete a paper diary to record their use of all products and also to answer questions on their cravings/urges and how satisfied they felt after using each product. Each participant receives a follow-up telephone call once the study is complete.

What are the possible benefits and risks of participating?

Participants will not gain any direct health benefits from taking part in this study. The screening examinations may help them learn about their general health. They may also help them discover an unknown medical condition. This study may help doctors, scientists or manufacturers learn things about tobacco products and e-cigarettes that will help people to quit smoking.

Participants can speak to the study doctor at any time to ask for advice on stopping using tobacco products. Nicotine and tobacco use can have side effects but participants included in the study will already be using tobacco products. They will likely not be exposed to levels higher than the ones they are usually exposed to. Tobacco products and e-cigarettes may cause side effects, such as cough, irritation in the mouth and throat, feeling faint, nausea, dizziness and headache. Less common side effects include nasal congestion, stomach discomfort, hiccups, vomiting, chest palpitations and cardiac arrhythmia. Taking blood and having a cannula inserted can be painful and may cause bleeding, bruising, discomfort, dizziness, infections and/or pain at the needle site. Some people may also feel faint or light-headed when blood is taken.

Where is the study run from?

Celerion (UK)

When is the study starting and how long is it expected to run for?

December 3rd, 2014 to December 20th, 2014.

Who is funding the study?

British American Tobacco (Investments) Limited (UK)

Who is the main contact?

Dr Brendan Colgan

brendan.colgan@celerion.com

## Contact information

**Type(s)**

Scientific

**Contact name**

Dr Brendan Colgan

**Contact details**

Celerion

22-24 Lisburn Road

Belfast

United Kingdom

BT9 6AD

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

## Secondary identifying numbers

BAT-OO1-PK/CA15374

# Study information

## Scientific Title

A part-randomised, part-blinded pharmacokinetic study of an e-cigarette containing different solutions and a combustible cigarette in healthy subjects

## Study hypothesis

Nicotine delivery and urge relief/satisfaction will be modified by the inclusion of different components/ingredients in the e-cigarette liquid

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Health and Social Care Research Ethics Committee B, 21/11/2014 ref: 14/NI/1118

## Study design

Single-centre part-randomised and part-blinded interventional study

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Other

## Study type(s)

Treatment

## Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

## Condition

Smoking

## Interventions

Subjects will be asked to either smoke a regular cigarette or use an e-cigarette containing one of three different solutions

## Intervention Type

Other

## Phase

Not Applicable

### **Primary outcome measure**

1. Plasma nicotine concentration C<sub>max</sub> and T<sub>max</sub>. Measured at timepoints -5, 1, 2, 3, 4, 5, 6, 7, 8, 10, 14.5, 30, 45, 60, 75 and 90 minutes following the start of product use
2. Subjective measurements of cravings/urges. Measured at timepoints -2, 1.25, 3.25, 5.5, 7.5, 10.5, 14.5, 30, 45, 60, 75 and 90 minutes following the start of product use. This will be done using a standard single-item question of cravings/urges.
3. Product satisfaction. This will be done using a customised panel examining numerous aspects of satisfaction with product use.

### **Secondary outcome measures**

1. Product (cigarette and e-cigarette) consumption per day during ambulatory periods
2. Heart rate during the first 12 minutes of product use
3. Cotinine and hydroxycotinine at the same timepoints as nicotine measurements

### **Overall study start date**

03/12/2014

### **Overall study end date**

07/08/2015

## **Eligibility**

### **Participant inclusion criteria**

1. Subjects will be males or non-pregnant, non-lactating females between 21 and 55 years of age inclusive
2. Women of child-bearing potential should be using one of the following acceptable methods of contraception: combined (estrogen and progestogen containing) oral, intravaginal or transdermal hormonal contraception associated with inhibition of ovulation; progestogen-only hormonal contraception, either oral, injected or implanted, associated with inhibition of ovulation; progestogen-only oral hormonal contraception where inhibition of ovulation is not the primary mode of action; intrauterine device (IUD); intrauterine hormone-releasing system (IUS); bilateral tubal occlusion; male or female condom with or without spermicide; cap, diaphragm or sponge with spermicide.
3. Women of nonchildbearing potential may be included if they are either surgically sterile (hysterectomy and/or oophorectomy) or postmenopausal for more than 1 year (with follicle-stimulating hormone [FSH] level greater than or equal to 8 mIU/mL) and must have a negative serum pregnancy test result during screening. Women who are surgically sterile must provide documentation of the procedure by an operative report.
4. Subjects must be in good health as determined by medical history, vital signs, blood biochemistry, haematology, urinalysis and physical examination.
5. Subjects with negative HIV and Hepatitis B and C results.
6. Subjects must have a body mass index (BMI) between 18 and 30 kg/m<sup>2</sup> inclusive. Male subjects must have a weight between 50 and 110kg and female subjects between 40 and 90kg.
7. No clinically significant abnormalities in blood pressure values (the differences between supine and standing BP are less than 20 mmHg) with no symptomatic evidence of postural hypotension.
8. Subjects will have negative results for the urinary drug of abuse and ethanol screening test.
9. Subjects will have given their written informed consent to participate in the study and to abide by the study restrictions.
10. Prior to study start, subjects must be daily smokers of at least 10 factory-produced

cigarettes and have smoked continuously for a minimum of one year. Subjects current brand of cigarette will have a machine-smoked ISO tar yield of 8-10 mg. Smoking status will be confirmed with a Smokerlyzer breath CO >10ppm and a urinary cotinine level of >200ng/ml at screening. Subjects should be familiar with the use of e cigarettes but not currently dual-using cigarettes and e-cigarettes

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

24

**Participant exclusion criteria**

1. Subjects who have a history of, or clinically active significant, neurological, gastrointestinal, renal, hepatic, cardiovascular, psychiatric, respiratory, metabolic, endocrine, haematological disease or other major disorders.
2. Subjects with significant allergies who in the opinion of the Principal Investigator should not be included.
3. Subjects who have been diagnosed with urticaria or asthma.
4. Subjects with a recent history of or current drug or alcohol abuse who in the opinion of the Investigator should not be included. Excessive intake of alcohol within the last 6 months, defined as a regular maximum weekly intake of greater than 7 drinks for women or 14 drinks for men. One drink is defined as one pint of regular beer (5% alcohol), 200 ml of wine (12% alcohol), or 25 ml of distilled spirits (40% alcohol).
5. Subjects with an inability to communicate well with the Investigator/study staff (i.e., language problem, poor mental development or impaired cerebral function).
6. Subjects who are participating in another clinical research study or who have participated in a clinical research study in the last 3 months.
7. Subjects who have had treatment with prescription medications within 21 days or over-the-counter medication within 72 hours of the planned first product use occasion. For female subjects, oral contraceptives, hormonal contraceptive devices and replacement hormonal therapies are not included in the list of drugs leading to exclusion.
8. Subjects who have used any drugs or substances (except tobacco) known to be strong inducers or inhibitors of any CYP enzymes (formerly known as cytochrome P450 enzymes) within a 28 days period prior to first product administration. For a list of such drugs and substances, please refer to <http://medicine.iupui.edu/clinpharm/ddis/main-table/>.
9. Subjects who have had any treatment with smoking cessation medications (e.g. Bupropion, Champix, any nicotine replacement therapy or nicotine delivery systems) within 30 days of the planned first product use occasion.
10. Subjects with any other clinically significant medical history, in the Investigator's opinion, including conditions which might affect drug absorption, metabolism or excretion.
11. Female subjects, who are pregnant or become pregnant during the course of the study.

12. Subjects who have lost or donated more than 450ml blood, plasma or platelets within the 3 months preceding the first product administration.
13. Subjects who are currently trying to stop smoking or considering stopping in the next two months.
14. Subjects who are unwilling or unable to comply with the study requirements.
15. Subjects who in the opinion of the Investigator should not participate in the study for any other reason.

**Recruitment start date**

03/12/2014

**Recruitment end date**

20/12/2014

## Locations

**Countries of recruitment**

Northern Ireland

United Kingdom

**Study participating centre**

**Celerion**

Belfast

United Kingdom

BT9 6AD

## Sponsor information

**Organisation**

British American Tobacco (Investments) Limited (UK)

**Sponsor details**

Group Research & Development

British American Tobacco

Regents Park Road

Southampton

United Kingdom

SO15 8TL

**Sponsor type**

Industry

**Website**

<http://www.bat-science.com>

ROR

<https://ror.org/01znsh139>

## Funder(s)

### Funder type

Industry

### Funder Name

British American Tobacco (Investments) Limited (UK)

## Results and Publications

### Publication and dissemination plan

This study will be written up for conference abstracts and a peer-reviewed publication along with a subsequent study that was run (NCT02474849).

### Intention to publish date

31/12/2016

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/01/2017		Yes	No
<a href="#">HRA research summary</a>			28/06/2023	No	No